

FINAL REPORT: Strategic Applications of Ultracold Atoms

Table of Contents

<i>Item</i>	<i>Page</i>
SF298	4
Statement of Problem	5
Summary of Most Important Results	6
Publications	9
Personnel	19
Report of Inventions	19

REPORT DOCUMENTATION PAGE			Form Approved OMB NO. 0704-0188
Public Reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comment regarding this burden estimates or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188,) Washington, DC 20503.			
1. AGENCY USE ONLY (Leave Blank)	2. REPORT DATE 5/20/04	3. REPORT TYPE AND DATES COVERED Final Report, FY00-03	
4. TITLE AND SUBTITLE Strategic Applications of Ultracold Atoms		5. FUNDING NUMBERS DAAD-19-00-1-0169	
6. AUTHOR(S) Mark Kasevich			
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Yale University, Sloane Physics Lab, 217 Prospect St., New Haven, CT 06520-8120		8. PERFORMING ORGANIZATION REPORT NUMBER 3 Yale University	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U. S. Army Research Office P.O. Box 12211 Research Triangle Park, NC 27709-2211		10. SPONSORING / MONITORING AGENCY REPORT NUMBER ARO <i>41064.1 - PH-MUR</i>	
11. SUPPLEMENTARY NOTES The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision, unless so designated by other documentation.			
12 a. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution unlimited.		12 b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) This consortium has initiated a focused collaborative program to advance matter wave sensors. We seek to combine atom interferometry with atom lasers and atom waveguides with the prospect of improving the sensitivity of such sensors by orders of magnitude as compared with existing state-of-the-art sensors. We will identify, explore and exploit fundamental scientific possibilities surrounding the production, manipulation and detection of ultra-cold atoms for a variety of sensing applications. Such sensors include gravimeters, gravity gradiometers, gyroscopes, magnetometers and frequency standards and have applications in science and technology and within the DoD. Sensitive and accurate inertial force sensors can be used in covert/passive navigation, precision guidance, underground structure detection, gravitational mapping, etc. They are non-emanating and capable of operating in a jammed-GPS environment. We seek to build awareness of DoD needs critical to national defense at the graduate training level, and to establish a dialogue between DoD and industrial researchers/managers and PhD trainees. The institutions identified in the proposed consortium attract talented students who are likely to become future leaders in science and technology.			
14. SUBJECT TERMS Atom Optics, Atom Interferometry, Bose-Einstein Condensation		15. NUMBER OF PAGES <i>15</i>	
		16. PRICE CODE	
17. SECURITY CLASSIFICATION OR REPORT UNCLASSIFIED	18. SECURITY CLASSIFICATION ON THIS PAGE UNCLASSIFIED	19. SECURITY CLASSIFICATION OF ABSTRACT UNCLASSIFIED	20. LIMITATION OF ABSTRACT UL

FINAL REPORT: Strategic Applications of Ultracold Atoms
Consortium members; Arizona, Harvard, MIT, Stanford, Yale

Statement of Problem Studied




Fig. 1. Atom laser gyro (ALG). An active atom laser gain element (blue) injects coherent de Broglie waves into 10 cm x 10 cm microfabricated ring resonator (red) and a planar substrate (black). Atom detection (yellow) is at the Heisenberg limit using quantum state manipulation techniques.

This consortium proposes a focused collaborative program to advance matter wave sensors. We want to combine atom interferometry with atom lasers and atom waveguides with the prospect of improving the sensitivity of such sensors by orders of magnitude as compared with existing state-of-the-art sensors. We will identify, explore and exploit fundamental scientific possibilities surrounding the production, manipulation and detection of ultra-cold atoms for a variety of sensing applications. Such sensors include gravimeters, gravity gradiometers, gyroscopes, magnetometers and frequency standards and have applications in science and technology and within the DoD.

Sensitive and accurate inertial force sensors can be used in covert/passive navigation, precision guidance, underground structure detection, gravitational mapping, *etc.* They are non-emitting and capable of operating in a jammed-GPS environment.

The recent demonstration of atom lasers and the recent progress in atom optics/interferometry raise the tantalizing prospect of a new class of ultra-sensitive sensors drawing on advances in both areas (Figs. 1 and 2). Compact, field-ready versions of these sensors could be orders of magnitude more sensitive than current state-of-the-art sensors.

As a concrete example of how developments in the atom laser and atom optics/interferometry fields will impact sensor technology, we consider the impact of advances in atom source, atom optics and atom readout on precision rotation sensing. Fig. 1 illustrates our vision of a prototype Sagnac effect rotation sensor, an atom-laser gyroscope. We want to emphasize that *all* components described below have already been demonstrated or are under development in at least one of the collaborating groups. The device operating principles are as follows: Single mode atom waveguides are used to guide atoms in a ring resonator configuration (analogous to the resonator of a ring laser gyroscope). An active atom gain element is inserted in the ring to inject de Broglie waves into the ring. Atoms are outcoupled at a waveguide beamsplitter and mixed on an atom-heterodyne detector. Squeezed state detection techniques are used to detect rotation shifts below the shot-noise limit. Spurious interferometer phase shifts, due to atom-atom interactions, are suppressed by tuning the atom-atom interaction strength to zero using Feshbach resonances. Intracavity loss from three-body collisions and output coupling of atoms is compensated for by the active gain element. To estimate the possible performance of this device, we assume that 10^6 atoms/sec can be coherently coupled into the ring through the active gain element and then detected at the Heisenberg-limit, that the ring has an area of 100 cm^2 (for a 10 cm x 10 cm device), and that an atom transits the loop an average of 10 times before being outcoupled. In this case, the device can resolve rotations of $2 \times 10^{-15} \text{ rad/sec}$ after 1 second. This is a factor of 10^6 improvement over the current state-of-the-art.

Minor modifications in the waveguide topologies allow for the realization of compact acceleration sensors such as gravimeters and gravity gradiometers, with similar gains in sensitivity expected. Assumptions similar to those of the previous paragraph lead to estimated accelerometer sensitivities of 10^{-14} g in 1 second and sensitivities to gravity gradients of 10^{-3} E ($1 \text{ E} = 10^{-9} \text{ sec}^{-2}$) after 1 second.




Fig 2. Interference fringes of two overlapping Bose-Einstein condensates [Error! Bookmark not defined.]. A major goal of the consortium is to develop this first demonstration into high-precision matter-wave sensors. This will be done by combining state-of-the art atom interferometry, BEC atom sources and atom wave guides.

Summary of most important results

Major program accomplishments in the three year performance interval include: 1) development of methods to create and detect entangled quantum states relevant to sub-shot noise interferometer read-out; 2) demonstration of Bose-condensate atom transport in waveguides; 3) demonstration of Bose-Einstein condensation on a micro-fabricated chip; 4) demonstration of novel waveguide structures for cold-atoms; 5) demonstration of a Bragg grating atom interferometer; 6) demonstration of a CW atom laser and 7) study of atom-surface interactions relevant to microfabricated waveguide devices. These experimental accomplishments, which have significantly advanced understanding of basic physical principles relevant to next generation atom interference sensors, have been supported by substantial advances in the understanding of degenerate atomic systems by the theory collaboration.

Accomplishments (organized by consortium performer):

Cold-atom guides (Prentiss; Harvard): Waveguide atom interferometers offer great potential for creating small, compact, robust rotation and acceleration sensors for guidance applications. Right now, magnetic field guiding is the leading candidate for realizing such guided interferometers. The Harvard group has developed waveguides based on ferromagnetic material, that allow very tight confinement to be obtained far above the physical surface of the magnetic structure, greatly reducing the heating of the atoms and even destruction of the atom sample that has resulted in experiments using atom waveguides based on current flowing through microfabricated wires. These structures have been used to produce atom clouds longer than 2 cm with more than 10^9 trapped atoms. A ferromagnetic ring waveguide has been designed and built, and cold atoms have been loaded into it.

BEC Interferometry (Prichard/Ketterle; MIT):

Progress in the field of atom optics depends on developing improved sources of matter waves and advances in their coherent manipulation. Miniaturizing the current carrying structures used to confine Bose-Einstein condensates offer prospects for finer control over the clouds. The MIT group has demonstrated that a gaseous Bose-Einstein condensate transported with optical tweezers can be transferred into a magnetic trap microfabricated on a silicon substrate. This has opened up a front on which further techniques for coherent condensate transport and manipulation can be explored. In their experiments, a condensate was released from the magnetic microtrap into a single-wire magnetic waveguide and its propagation characteristics were observed. Condensates were observed to propagate 12 mm before exiting the field-of-view of the imaging system. A single-mode (excitation-less) condensate propagation along homogeneous segments of the waveguide was observed. Transverse excitations were created in condensates propagating through perturbations in the guiding potential. These perturbations resulted from geometric deformations of the current carrying wires on the substrate. Finer imperfections were observed when trapped condensates were brought closer to the microchip as evidenced by the longitudinal fragmentation of the cloud. Such imperfections have to be controlled in order to use atom chips for precision atom interferometry.

Unlike other macroscopic quantum systems such as superfluid ^4He and optical lasers, dilute gas Bose-Einstein Condensates have so far been only produced in a pulsed mode. The MIT group has realized a continuous BEC source by periodically replenishing a condensate held in an optical dipole trap with new condensates. A moving optical tweezers for Bose-Einstein condensates was used to transport condensates from where they were produced into a reservoir optical trap. The freshly produced condensates periodically replenished the condensate in the reservoir trap, thereby continuously maintaining a condensate of more than 10^6 atoms. The crucial step in realizing a continuous BEC source was to make sure that the new cooling cycle did not destroy the condensate held in the reservoir trap. This involved shielding it from light during laser cooling, keeping it far away from the incoming hot atoms, and to hold it in an optical trap which made it immune against stray magnetic fields which were created during the evaporative cooling phase.

By colliding two Bose-Einstein condensates the MIT group has observed strong bosonic stimulation of the elastic scattering process. When a weak input beam (third wave) was applied as a seed, it was amplified by a factor of 20, and an initially unpopulated conjugate wave was created. This large gain atomic four-wave mixing resulted in the generation of two macroscopically occupied pair-correlated atomic beams. Since each collision process adds one atom each to the seed and conjugate waves, fluctuations in

the relative atom number are suppressed (squeezed). For the observed gain of twenty, the number fluctuations should be below the shot noise by a factor of $40^{1/2}$. Limitations for using collisions to create twin beams have been identified. These include loss by subsequent collisions and competition between other modes with similar gain.

Finally, the MIT group has demonstrated a new atom interferometer scheme, which shows promise for a high precision measurement of the recoil energy of an atom. A precise measurement of the recoil frequency will lead to a more precise determination of h/m and of the fine structure constant α . The interferometer extends previous schemes used at Stanford and NYU, and combines their advantages. Optical standing wave pulses were used to create a symmetric three-path interferometer. This configuration encodes the photon recoil phase in the contrast of the interference fringes, rather than in their phase. Because it is insensitive to the fringe phase, the method is not sensitive to vibrations, accelerations, or rotations. The symmetry also suppresses errors from magnetic field gradients, and our use of only one internal state suppresses errors arising from differences in the ac Stark shifts between different internal states. A crucial aspect of this new interferometer is the use of atomic samples with sub-recoil momentum distribution, in this case a BEC. This allows the contrast oscillations to persist for many cycles, permitting precise determination of the recoil phase in a single “shot.”

Waveguide BEC (Chu, Vuletic; Stanford): Bose-Einstein condensation has recently been achieved in a novel microchip topology. The experimental method to obtain BEC is robust and simple; it uses only a single vacuum chamber and a magnetic microtrap which is directly loaded from a standard magneto-optical trap. The short evaporation time of 3s is limited only by the axial trap vibration frequency, and it is estimated that the evaporation time can be shortened to 1s at a MOT loading time of 500ms. The condensates containing a few thousand rubidium atoms are produced at a distance of 60 μ m from the silicon substrate, and are stable to distances down to 15 μ m. An atom number shot noise-limited imaging system for the condensate has been set-up to study the condensate. In the near future, we will try to demonstrate sub-shot noise fluctuations (“squeezing”) in the relative atom number contained in different spatial regions of the condensate. This is equivalent to a determination of the average condensate position with sub-shot noise accuracy, and may be of use in simple interferometric schemes with condensates.

Squeezing and quantum state control in optical lattice (Kasevich; Yale/Stanford): The Yale/Stanford group demonstrated that the optical lattice system can be used to produce squeezed atomic states. The basic idea of this work is to exploit mean-field interactions to produce non-trivial many-body states of weakly connected ensembles of Bose condensates. Experimentally, arrays of condensates are confined in corrugated potentials in which atoms may resonantly tunnel between adjacent wells. This opens up the possibility of producing non-trivial many-body states. In particular, the ground state of the system can no longer be described by a factorizable quantum wavefunction (ie. as the simple product of identical single-particle wavefunctions). The two-well version of this problem is exactly solvable. In the limit where the mean-field interaction dominates tunneling, the total energy of the system is minimized by suppressing number fluctuations at each trapping site – that is by squeezing the on-site number distribution. The corresponding phase spread, via the number-phase uncertainty principle, increases. On the other hand, when tunneling dominates, the ground state approaches that many-body state associated with a non-interacting gas. In this case, the number fluctuations at a given site approach those of a coherent state. These effects have been demonstrated experimentally by analyzing the phase coherence of atoms in the lattice array. This work has recently been extended to study the Mott-insulating regime, both interferometrically and by transport measurements. This work has also identified several new paths to the creation of sub-shot noise interferometers.

Theory (Meystre, Glauber, Wright; Tuscon/Harvard): The theory collaboration has significantly advanced the state-of-knowledge for the quantum field theory of degenerate bosonic and fermionic systems, with particular emphasis to sensors applications. Most recently, the focus on fermionic systems may identify mitigation strategies to overcome some of the technology hurdles identified in the study of guided atom systems.

For example, it is only recently that it has been convincingly argued that atomic four-wave mixing is possible with fermions. However, the theory so far has been limited to an incident beam consisting of a single test particle. It also neglected the dynamics of the density grating. These limitations have now been removed by taking into account both a multi-particle beam and the back-action of the incident beam on the

fermionic density grating. Using this formalism, it has been shown that when the number of atoms in the incident beam becomes a sizable fraction of the number of atoms forming the density grating, it no longer decays away due to the dephasing resulting from the slightly different energies of the fermions forming the incident beam. Instead, it exhibits large nonlinear amplitude oscillations that are coupled to the Bragg oscillations of the beam. This leads to the efficient generation of the fourth scattered wave, even for times much longer than the grating dephasing time. It is well known that four-wave mixing with bosons can lead to the generation of squeezed states. Therefore a central feature of future work will examine the statistical properties of the scattered beam generated in fermionic four-wave mixing to see if it is possible to generate novel fermionic states. Phase conjugation should also be possible if one uses a superfluid Fermi gas due to the presence of anomalous moments resulting from the formation of Cooper pairs.

In related work, the phase resolution limit of a Mach-Zehnder atom interferometer whose input consists of degenerate quantum gases of either bosons or fermions has been analyzed. For degenerate gases, the number of atoms within one de Broglie wavelength is larger than unity, so that atom-atom interactions and quantum statistics are no longer negligible. It has been shown that for equal atom numbers, the phase resolution achievable with fermions can be noticeably better than for interacting bosons. This is a strong argument for a further extensive study of fermionic atom optics.

The recent experimental success in creating quantum-degenerate atomic Fermi gases is opening up fascinating new opportunities to explore the quantum statistics of ultracold atoms. Fermionic behavior is strongly constrained by the Pauli Exclusion Principle. This limits the variety of possible nonlinear atom optics effects, but also offers the potential for novel applications without analogs in optics. These include, for example, low-noise inertial and rotation sensors, and quantum information processing. In a new development of the theory research program, the new situation where a gas of bosons serves as a nonlinear medium for fermionic atoms has been explored. In particular, interatomic interaction between a Bose-Einstein condensate and a fermionic beam can be employed to manipulate the quantum state of the beam. As a first step, and drawing on an analogy to nonlinear optics, the interaction can be described in terms of an effective attractive Kerr nonlinearity, and show that a two-fermion bound state can result with a unique signature in a nonlinear atom optical experiment. Future study of this coupled Bose-Fermi systems may be of particular relevance for the manipulation of the quantum statistical properties of fermionic atomic beams, e.g. changes from antibunched and bunched beams, dynamic Cooper pairing, and the formation of quantum solitons in ultracold fermionic atomic beams.

Publications

Paper published in peer-reviewed journals:

- P. Meystre, Physics World **13**, 24 (2000).
- E. V. Goldstein, M. G. Moore and P. Meystre, Laser Physics **10**, 8 (2000).
- J. Heurich, M. G. Moore and P. Meystre, Optics Commun. **179**, 549 (2000).
- E. V. Goldstein, M. G. Moore and P. Meystre, in Advances in Laser Physics, edited by V. S. Letokhov and P. Meystre, Laser Science and Technology Second Series Volume 1, Harwood Academic Publishers, Amsterdam (2000), p. 117.
- G. Prataiviera, J. Zapata and P. Meystre, Phys. Rev. A **62**, 023605 (2000).
- H. Pu and P. Meystre, Phys. Rev. Lett. **85**, 3987 (2000).
- S. Poetting, E. S. Lee, W. Schmitt, I. Romyantsev, B. Mohring and P. Meystre, Phys. Rev. A **62**, 060101 (R)(2000).
- S. Poetting, O. Zobay, P. Meystre and E. M. Wright, J. Mod. Optics **47**, 2653 (2000).
- M. G. Moore and P. Meystre, Phys. Rev. Lett. **85**, 5026 (2000).
- E. V. Goldstein, M. G. Moore, H. Pu and P. Meystre, Phys. Rev. Lett. **85**, 5030 (2000).
- E. M. Wright, J. Arlt and K. Dholakia, Phys. Rev. A **63**, 013608 (2001).
- S. Inouye, R.F.Löw, S. Gupta, T. Pfau, A. Görlitz, T. L. Gustavson, D. E. Pritchard and W. Ketterle, Phys. Rev. Lett. **85**, 4225-4228 (2000).
- R. Onofrio, C. Raman, J. M. Vogels, J. Abo-Shaeer, A.P. Chikkatur, and W. Ketterle, Phys. Rev. Lett. **85**, 2228-2231 (2000).
- A.P. Chikkatur, A. Görlitz, D.M. Stamper-Kurn, S. Inouye, S. Gupta, and W. Ketterle, Phys. Rev. Lett. **85**, 483-486 (2000).
- J. McGuirk, M. Snadden and M. Kasevich, Phys. Rev. Lett. **85**, 4413 (2000).
- C. Chin, V. Vuletic, A.J. Kerman, and S. Chu, Phys. Rev. Lett. **85**, 2717-2720 (2000).
- C. Raman, J.R. Abo-Shaeer, J.M. Vogels, K. Xu, and W. Ketterle: Vortex Nucleation in a Stirred Bose-Einstein Condensate. Phys. Rev. Lett. **87**, 210402 (2001).
- A. Görlitz, J.M. Vogels, A.E. Leanhardt, C. Raman, T.L. Gustavson, J.R. Abo-Shaeer, A.P. Chikkatur, S. Gupta, S. Inouye, T. Rosenband, and W. Ketterle:
Realization of Bose-Einstein condensates in lower dimensions. Phys. Rev. Lett. **87**, 130402 (2001).
Squeezed States in a Bose-Einstein Condensate, C. Orzel, A. Tuchman, M. Fenselau, M. Yasuda, and M. Kasevich, Science **291**, 2386 (2001).
44. Low-noise Detection of Ultra-cold Atoms, J. McGuirk, G. Foster, J. Fixer and M. Kasevich, Opt. Lett. **26**, 364 (2001).
- S. Inouye, S. Gupta, T. Rosenband, A.P. Chikkatur, A. Görlitz, T.L. Gustavson, A.E. Leanhardt, D.E. Pritchard, and W. Ketterle: Observation of vortex phase singularities in Bose-Einstein condensates. Phys. Rev. Lett. **87**, 080402 (2001).

- J.R. Abo-Shaeer, C. Raman, J.M. Vogels, and W. Ketterle: Observation of Vortex Lattices in Bose-Einstein Condensates. *Science* 292, 476-479 (2001).
- Three-dimensional cavity Doppler cooling and cavity sideband cooling by coherent scattering, V. Vuletic, H.W. Chan, and A.T. Black, *Phys. Rev. A* 64, 033405 (2001).
- W. Ketterle and S. Inouye: Collective enhancement and suppression in Bose-Einstein condensates. *Compte rendus de l'académie des sciences, Série IV - Physique Astrophysique*, vol. 2, pp. 339-380 (2001); e-print cond-mat/0101424.
- C. Raman, R. Onofrio, J.M. Vogels, J.R. Abo-Shaeer, and W. Ketterle: Dissipationless flow and superfluidity in gaseous Bose-Einstein condensates. *J. Low Temp. Phys.* 122, 99-116 (2001).
- W. Ketterle and S. Inouye: Does matter wave amplification work for fermions? *Phys. Rev. Lett.* 86, 4203-4206 (2001). A. Görlitz, A.P. Chikkatur, and W. Ketterle: Enhancement and suppression of spontaneous emission and light scattering by quantum degeneracy. *Phys. Rev. A* 64, 041601(R) (2001).
- M. G. Moore and P. Meystre, "Parametric amplification of coupled atomic and optical fields", in "Directions in Quantum Optics", a volume in the Memory of D. F. Walls, edited by H. J. Carmichael, R. J. Glauber and M. O. Scully, Springer Verlag, Berlin, p. 116 (2001).
- J. Heurich, H. Pu, M. G. Moore, and P. Meystre, "Instabilities and self-oscillations in atomic four-wave mixing," *Phys. Rev. A* 63, 033605 (2001).
- M. G. Moore and P. Meystre, "Atomic four-wave mixing: fermions vs. bosons", *Phys. Rev. Lett.* 86, 4199 (2001).
- J. Zapata, A. M. Guzman, M. G. Moore, and P. Meystre, "Gravity-induced Wannier-Stark ladder in an optical lattice", *Phys. Rev. A* 63, 023607 (2001).
- P. Meystre, "Nonlinear optics of de Broglie waves", in "Advances in Atomic, Molecular and Optical Physics", Vol. 47, edited by B. Bederson and H. Walther, Academic Press, San Diego, p. 1, (2001).
- S. Raghavan, H. Pu, P. Meystre, and N. P. Bigelow, "Generation of arbitrary Dicke states in spinor Bose-Einstein condensates", *Optics Commun.* 188, 149 (2001).
- H. Pu, M. G. Moore, and P. Meystre, "Atomic squeezing and entanglement from Bose-Einstein condensates", in "Modern Challenges in Quantum Optics", edited by M. Orszag and J. C. Retamal, Springer Verlag (Berlin), p. 161 (2001).
- S. Pötting, M. Cramer, C. H. Schwalb, H. Pu, and P. Meystre, "Coherent acceleration of Bose-Einstein condensates", *Phys. Rev. A* 64, 023604 (2001).
- S. Pötting, M. Cramer, and P. Meystre, Momentum state engineering and control in Bose-Einstein condensates, *Phys. Rev. A* 64, 063613 (2001).
- H. Pu, W. Zhang, and P. Meystre, "Ferromagnetism in a lattice of Bose condensates", *Phys. Rev. Lett.* 87, 140405 (2001).
- P. Meystre, "Atom Optics", Springer-Verlag, New York (2001).
- M. D. Girardeau, E. M. Wright, and J. M. Triscari, "Ground state properties of a one-dimensional condensate of hard core bosons in a harmonic trap," *Phys. Rev. A* 63, 033601 (2001).

- M. D. Girardeau and E. M. Wright, "Measurement of one-particle correlations and momentum distributions for trapped 1D gases," *Phys. Rev. Lett.* 87, 050403 (2001).
- J. Arlt, K. Dholakia, J. Soneson, and E. M. Wright, "Optical dipole traps and atomic waveguides based on Bessel light beams," *Phys. Rev. A* 63, 63602 (2001).
- M. D. Girardeau and E. M. Wright, "Bose-Fermi variational theory of the Bose-Einstein condensate crossover to the Tonks gas," *Phys. Rev. Lett.* 87, 210401 (2001).
- M. D. Girardeau and E. M. Wright, "Quantum mechanics of one-dimensional trapped Tonks gases," *Laser Physics* 12, 8 (2002).
- P. Anderson and P. Meystre, "Nonlinear atom optics," *Contemporary Physics* 44, 473 (2003).
- Chris P. Search, Weiping Zhang, and P. Meystre, "A molecular micromaser," *Phys. Rev. Lett.* 91, 190401 (2003).
- Han Pu, Weiping Zhang, and Pierre Meystre, "Wave mixing of optical pulses and Bose-Einstein condensates," *Phys. Rev. Lett.* 91, 150407 (2003).
- P. Search and Pierre Meystre, "Phase conjugation of a quantum-degenerate atomic Fermi beam," *Phys. Rev. Lett.* 91, 100401 (2003).
- Weiping Zhang, Ewan M. Wright, Han Pu, and Pierre Meystre, "A fundamental limit for integrated atom optics with Bose-Einstein condensates," *Phys. Rev. A* 68, 023605 (2003).
- H. Christ, C. P. Search, and P. Meystre, "Dynamics of fermionic four-wave mixing," *Phys. Rev. A* 67, 063611 (2003).
- T. Miyakawa, H. Christ, C. P. Search, and P. Meystre, "Four-wave mixing in degenerate Fermi gases: Beyond the undepleted pump approximation," *Phys. Rev. A* 67, 063603 (2003).
- P. Search and P. Meystre, "Noise limits in matter-wave interferometry using degenerate quantum gases," *Phys. Rev. A (Rapid Communication)* 67, 061601 (2003).
- W. Zhang, H. Pu, C. P. Search, P. Meystre, and E. M. Wright, "Two-fermion bound states in a Bose-Einstein condensate," *Phys. Rev. A (Rapid Communication)* 67, 021601(R) (2003).
- S. Pötting, P. Meystre and E. M. Wright, "Atomic solitons in optical lattices", in "Nonlinear Photonic Crystals," edited by R. L. Slusher and B. Eggleton, Springer Verlag (2003).
- W. Zhang, Chris P. Search, Han Pu, Pierre Meystre, and Ewan M. Wright, "Feshbach resonance-induced atomic filamentation and quantum pair correlation in atom-laser-beam propagation," *Phys. Rev. Lett.* 90, 140401 (2003).
- H. Pu, L. O. Baksmaty, W. Zhang, N. P. Bigelow, and P. Meystre, "Effective-mass analysis of Bose-Einstein condensates in optical lattices: Stabilization and levitation," *Phys. Rev. A* 67, 043605 (2003).
- B. P. Anderson, K. Dholakia, and E. M. Wright, "Atomic-phase interference devices based on ring-shaped Bose-Einstein condensates: Two ring case," *Phys. Rev. A* 67 033601 (2003).
- P. Meystre, "Atom Optics", Japanese Translation, Springer Verlag, Tokyo (2003).
- "Suppression of photon rescattering due to spatial anisotropy in a cold atomic gas", M. Vengalattore, W. Rooijakkers, R. Conroy, and M. Prentiss, *Phys. Rev. A* 67, 063412 (June 2003).

“Compact, robust source of cold atoms for efficient loading of a magnetic guide”, R.S. Conroy, Y. Xiao, M. Vengalattore, W. Rooijakkers, and M. Prentiss, Optics Commun. 226, 259-266 (October 2003).

“Observation of Caustics in the Trajectories of Cold Atoms in a Linear Magnetic Potential”, W. Rooijakkers, S. Wu, P. Striehl, M. Vengalattore and M. Prentiss, Phys. Rev. A 68, 063412 (December 2003).

A. E. Leanhardt, T. A. Pasquini, M. Saba, A. Schirotzek, Y. Shin, D. Kielpinski, D. E. Pritchard, and W. Ketterle: Cooling of Bose-Einstein condensates below 500 Picokelvin. Science 301, 1513-1515 (2003).

Y. Shin, M. Saba, T. Pasquini, W. Ketterle, D.E. Pritchard, and A.E. Leanhardt:

Atom interferometry with Bose-Einstein condensates in a double-well potential, Phys. Rev. Lett., in print; cond-mat/0306305.

Observation of Collective Friction Forces due to Self-Organization of Atoms: From Rayleigh to Bragg Scattering, A.T. Black, H.W. Chan, and V. Vuletic, Phys. Rev. Lett. 91, 203001 (1-4) (2003).

Papers published in non-peer-reviewed journals:

V. Vuletic, A.J. Kerman, C. Chin, and S. Chu (Florence, June 2000), published in Proceedings of the XVII. International Conference on Atomic Physics (ICAP 2000), edited by P. De Natale (American Institute of Physics).

A.J. Kerman, C. Chin, V. Vuletic, S. Chu, Paul J. Leo, Carl J. Williams, and P.S. Julienne, in Proceedings of the Euroconference on Atomic Optics and Interferometry (Corsica, July 2000).

Atom Interferometry with Cold Atoms, M. Kasevich, Enrice Physics School Lectures, in press, 2000.

Experimental Studies of Bose-Einstein Condensates in Sodium. in: Bose Einstein Condensates and Atom Lasers, edited by S. Martellucci, A.N. Chester, A. Aspect, and M. Inguscio (Kluwer Academic, New York, 2000), pp. 1-29.

A.J. Kerman, C. Chin, V. Vuletic, S. Chu, Paul J. Leo, Carl J. Williams, and P.S. Julienne, published in Proceedings of the Euroconference on Atomic Optics and Interferometry (Corsica, July 2000).

Cavity cooling with a hot cavity, V. Vuletic, in Laser Physics at the Limits, edited by H. Figger, D. Meschede, and C. Zimmermann, pp. 67-74 (Springer 2001);

Coherent Manipulation of Cold Atoms in Optical Lattices for a Possible Quantum Computation System, C. Chin, V. Vuletic, A.J. Kerman, and S. Chu, in Proceedings of the 15th International Conference on Laser Spectroscopy (ICOLS 2001, Snowbird), edited by S. Chu, V. Vuletic, A.J. Kerman, and C. Chin (World Scientific, Singapore 2001).

Paper presented at meetings, but not published in conference proceedings:

A. Görlitz, A.P. Chikkatur, S. Inouye, S. Gupta, T. Pfau, D.M. Stamper-Kurn, D.E. Pritchard, and W. Ketterle, 16th Interdisciplinary Laser Science Conference (ILS-XVI), Providence, RI, October 22-26, 2000, Conference Program, p. 96.

S. Inouye, R. Löw, T. Pfau, S. Gupta, A.P. Chikkatur, A. Görlitz, D.E. Pritchard, and W. Ketterle, 16th Interdisciplinary Laser Science Conference (ILS-XVI), Providence, RI, October 22-26, 2000, Conference Program, p. 96.

W. Ketterle, 2000 Atomic, Molecular and Optical Physics Research Meeting, U.S. Department of Energy, Office of Basic Energy Sciences, Warrenton, Virginia, September 26-29, 2000, Book of Abstracts.

T.L. Gustavson, J. Abo-Shaeer, A.P. Chikkatur, A. Görlitz, S. Gupta, Z. Hadzibabic, S. Inouye, R. Löw, R. Onofrio, C. Raman, J.M. Vogels, D.E. Pritchard, and W. Ketterle, 6th Workshop on Atom Optics and Interferometry, Cargese, Corse (France), July 26-29, 2000, Invited Talks Book of Abstracts.

S. Gupta, A.P. Chikkatur, A. Görlitz, T.L. Gustavson, S. Inouye, A.E. Leanhardt, R. Löw, T. Pfau, T. Rosenband, D.E. Pritchard, and W. Ketterle, 6th Workshop on Atom Optics and Interferometry, Cargese, Corse (France), July 26-29, 2000, Book of Poster Abstracts.

Z. Hadzibabic, J.R. Abo-Shaeer, D.S. Durfee, J. Gore, M. Köhl, C.E. Kuklewicz, R. Onofrio, C. Raman, J.M. Vogels, and W. Ketterle, 6th Workshop on Atom Optics and Interferometry, Cargese, Corse (France), July 26-29, 2000, Book of Poster Abstracts.

J.M. Vogels and W. Ketterle, Workshop on Rotating Bose-Einstein condensates, Trento, June 11-13, 2000, Book of abstracts p. 13.

W. Ketterle, ICAP 2000, Florence, June 4-9, 2000, Book of Abstracts, p. 47.

V. Vuletic, C. Chin, A.J. Kerman, and S. Chu, Quantum Electronics and Laser Science Conference (San Francisco, May 2000).

V. Vuletic, A.J. Kerman, C. Chin, and S. Chu (Florence, June 2000), published in Proceedings of the XVII. International Conference on Atomic Physics (ICAP 2000), edited by P. De Natale (American Institute of Physics).

E. M. Wright, M. D. Girardeau, and J. M. Triscari, "Exact many-body solutions for one dimensional ultracold trapped gases," Los Alamos Days Meeting, Tucson (Feb. 2001)

E. M. Wright and M. D. Girardeau, "Quantum mechanics of 1D trapped Tonks gases," ITAMP, Harvard-Smithsonian, Cambridge (March 2001).

E. M. Wright, J. Arlt, and K. Dholakia, "Toroidal optical dipole traps for two dimensional BECs," QELS 2001, Baltimore, MD (May 2001).

M. D. Girardeau and E. M. Wright, "Theory of de Broglie atomic waveguides," Prokhorov Fest, Moscow (June 2001).

E. M. Wright, "Schizophrenic atoms in one-dimension," Research Colloquium, University of St. Andrews (July 2001).

E. M. Wright, "Quantum mechanics of 1D trapped Tonks gases," Research Colloquium, University of Strathclyde (July 2001).

P. Meystre, "Nonlinear atom optics: bosons vs. fermions", 8th Rochester Conference on Coherence and Quantum Optics, Rochester, NY (2001).

P. Meystre, "Nonlinear atom optics: Recent results and prospects", Plenary talk, International Laser Physics Conference PHYS'01, Moscow, Russia (2001).

P. Meystre, "Nonlinear atom optics of bosons and fermions", Gordon Research Conference on Nonlinear Optics, New London, NH (2001).

P. Meystre, "Nonlinear atom optics --- From de Broglie to Heisenberg", Alexander von Humboldt 18th Symposium, "100 Years Werner Heisenberg --- Works and Impact", Bamberg, Germany (2001).

P. Meystre, "Nonlinear atom optics", Symposium in the Honor of Professors Chatelain and Quattropani", Swiss Federal Institute of Technology, Lausanne, Switzerland (2001).

M. Cramer, C. Schwalb, S. Poetting, H. Pu and P. Meystre, "Coherent acceleration of Bose-Einstein condensates", 2nd Pan-Pacific Basin Workshop on Microgravity Sciences, Pasadena, CA (2001).

Sierk Poetting, Marcus Cramer, Christian H. Schwalb, Han Pu, and Pierre Meystre, "Lossless acceleration of Bose-Einstein condensates", DAMOP Annual Meeting, London, Ontario (2001).

H. Pu, W. Zhang and P. Meystre, "Phonon excitations and stability of a Bose-Fermi mixture quantum gas", 8th Rochester Conference on Coherence and Quantum Optics, Rochester, NY (2001).

Cold atoms in anisotropic traps and in colored vacua, Seminar, Joint Atomic Physics series, Harvard University (2001).

Cavity Doppler cooling in three dimensions, V. Vuletic, A. T. Black, and H. W. Chan, Quantum Electronics and Laser Science Conference (QELS) 2001, (Baltimore 2001), invited presentation.

Cavity cooling: Towards laser cooling of light scatterers? V. Vuletic, H. Chan, and A. Black, Gordon Research Conference on Atomic Physics, (Williamstown 2001), invited presentation.

Three-dimensional cavity Doppler cooling by coherent scattering, V. Vuletic, A. Black, and H. Chan, International Workshop on Quantum Gases, (Konstanz 2001), invited presentation.

Collective enhancement and suppression in Bose-Einstein condensates.
Winter Colloquium on the Physics of Quantum Electronics, Snowbird, Utah,
1/8-11

Japan workshop on cold atoms
(Talk by A. Chikkatur)

Amplification of Atoms and Light in Bose-Einstein condensates.
plenary talk, APS April Meeting, Washington, D.C., 4/28 – 5/1, 2001

Experiments with a Bose-Einstein Condensate in an Optical Trap.
DAMOP 2001, London, Ontario, Canada, 5/16 – 5/19, 2001.
(Talk by T. Gustavson)

Vortex Matter in Bose-Einstein Condensates.
International Conference on Squeezed states and Uncertainty Relations, Boston,
MA, June 4 – 8 (ICSSUR 2001)
(Talk by Chandra Raman)

Optical trapping and manipulation of Bose-Einstein condensates.
15th International Conference on Laser Spectroscopy, Snowbird, Utah, June 10-15,
2001 (ICOLS 01
(Talk by T. Gustavson)

Coherence and collective behavior in Bose-Einstein condensates.
plenary talk, Eight Rochester Conference on Coherence and Quantum Optics,

Rochester, June 13 -16.

Spinning and shaping condensates with laser beams.
Atomic Physics Gordon conference, Williamstown, MA, June 17 – 22.

Bose-Einstein condensation – quantum mechanics near zero temperature.
plenary talk, Conference on statistical physics STATPHYS 21, Cancún, Mexico, July 18.

Recent BEC experiments at MIT – BEC in large-volume optical traps and interference of vortices. International Workshop “Quantum Gases”, Insel Reichenau, Germany, July 19 –July 21, 2001.

Vortex Excitations in a Bose-Einstein Condensate.
International Symposium on Foundations of Quantum Mechanics (ISQM-Tokyo '01), Hatoyama, Japan, 8/27-30.
(Talk by Shin Inouye)

Optical Manipulation of Bose-Einstein Condensates.
"Bose-Einstein Condensation", EuroConference on the Physics of Atomic Gases at Low Temperatures, San Feliu de Guixols, Spain, September 15-20.
(Talk by A. Görlitz)

Vortex Lattices in Bose-Einstein Condensates.
"Bose-Einstein Condensation", EuroConference on the Physics of Atomic Gases at Low Temperatures, San Feliu de Guixols, Spain, September 15-20.
(Talk by J. Abo-Shaeer)

Ketterle, Superfluidity and Coherence in Bose-Einstein Condensates.
Nobel Jubilee Symposium, Gothenborg, Sweden, December 4.

Ketterle, When atoms behave as waves: Bose-Einstein condensation and the atom laser.
Nobel Lecture, Stockholm, Sweden, December 8.

Kasevich, ACOLS 2002, Plenary, Brisbane, Dec. 2001.

Kasevich, BEC Symposium, Kiola, Australia, Dec. 2001.

Kasevich, ITAMP Symposium, Boston, Nov. 2001.

Kasevich, University of Maryland, Physics Dept. Colloquium, Oct. 2001.

Kasevich, FSM, St. Andrews, Sept. 2001.

Kasevich, ICOLS, Utah, June 2001.

Kasevich, PMOSSP Seminar (Navy), Mitchel Field, May 2001.

Kasevich, ONR Nav. Tech. Review, Washington D.C., May 2001.

Kasevich, NTAG-2001, BIPM, Paris, Feb. 2001 (talk given by G. Foster)

Kasevich, Laser Cooling and Applications, Japan, Jan. 2001 (talk given by Jeff McGuirk).

"Atom optics," Lecture Series (10 hours), Troisième Cycle Romand de Physique, Swiss Federal Institute of Technology, Lausanne, Switzerland (2003).

“Fermionic atom optics --- a tutorial,” invited talk, TAMU-ONR-DARPA Workshop on Quantum Optics, Grand Targhee, Wyoming (2003).

“Cavity de Broglie optics --- a molecular micromaser,” invited talk, QUEST 2003 Symposium, Santa Fe, New Mexico (2003).

“Four-wave mixing of fermionic matter waves,” invited talk, 12th International Laser Physics Workshop LPHYS'03, Hamburg, Germany (2003).

“Nonlinear mixing of pulsed optical and matter waves,” invited talk, 12th International Laser Physics Workshop LPHYS'03, Hamburg, Germany (2003).

“Cavity de Broglie optics - The molecular micromaser,” invited talk, 2003 US-Japan Joint Seminar ‘Quantum Correlations and Coherence’, Yatsugatake, Japan (2003).

“Quantum pair correlation in the Feshbach-resonance-induced atomic filamentation of a propagating atom-laser beam,” Weiping Zhang, C. P. Search, Han Pu, P. Meystre, and E. M. Wright, APS Division of Atomic, Molecular and Optical Physics Annual Meeting, Boulder (CO), 2003.

“Noise limits in matter-wave interferometry using degenerate quantum gases,” C. P. Search and P. Meystre, APS Division of Atomic, Molecular and Optical Physics Annual Meeting, Boulder (CO), 2003.

Bose-Einstein Condensates in Magnetic Waveguides.
The 5th Laser Cooling Workshop, Awaji Yumebutai, Japan, 1/7/2003.
(Talk by Aaron Leanhardt).

Ultrakalte Quantengase – die kälteste Materie im Universum.
Plenary Talk, Spring Meeting of the German Physical Society (DPG), Hannover, 3/26/2003.

Small Scales and Short Times: Two Recent Experiments with Bose-Einstein Condensates.
International Workshop Quantum Gases, Schloß Reinsburg, Günzburg, Germany, 4/2 – 4/5/2003.
(Talk by Dominik Schneble).

Quantum Effects on a Macroscopic Scale: BEC in Dilute Atomic Gases.
Symposium in honor of Prof. G. Leung, Physics Department, University of Massachusetts, North Dartmouth, MA, 4/11 – 4/12/2003).
(Talk by Dominik Schneble).

Single Mode Atom Chips for BECs.
Gordon Conference on Atomic Physics, Tilton, NH, 6/19/2003.
(Talk by D.E. Pritchard).

Recent results at MIT on cold bosons and fermions.
Bose-Einstein Condensation, EuroConference on the New Trends in Physics of Quantum Gases, San Feliu de Guixols, Spain, 9/13 – 9/18/2003.

Collective cavity cooling of cesium atoms, H.W. Chan, A.T. Black, and V. Vuletic, Physics Colloquium, NYU (New York 2003).

Bose-Einstein condensates near room-temperature surfaces Casimir potential and Johnson noise, Y. Lin, I. Teper, C. Chin, and V. Vuletic, DAMOP meeting (Boulder 2003).

Collective Friction Forces due to Self-Organization of Cesium Atoms, A.T. Black, H.W. Chan, and V. Vuletic, 16th International Conference on Laser Spectroscopy (ICOLS 2003, Palm Cove, Australia).

Collective Friction Forces due to Self-Organization of Cesium Atoms: From Rayleigh to Bragg Scattering, A.T. Black, H.W. Chan, and V. Vuletic, 9th Japan-US Joint Seminar on Quantum Correlation and Coherence (Yatsugatake, Japan 2003).

Stability of Bose-Einstein condensates near room-temperature surfaces, Y. Lin, I. Teper, C. Chin, and V. Vuletic, Physics Colloquium, York University (Toronto 2003).

Collective laser cooling due to spatial self-organization of classical atoms: From Rayleigh to Bragg scattering, A.T. Black, H.W. Chan, and V. Vuletic, Physics Colloquium, University of Toronto (2003).

Collective Friction Forces due to Spatial Self-Organization of Atoms: From Rayleigh to Bragg Scattering, A.T. Black, J. Thompson, H.W. Chan, and V. Vuletic Joint Atomic Physics Series, Harvard University (Cambridge 2004).

Manuscripts submitted, but not yet published:

Chris P. Search, Weiping Zhang, and Pierre Meystre, "Inhibiting three-body recombination in atomic Bose-Einstein condensates," to be published in Phys. Rev. Lett.

D. Meiser, C. P. Search, and P. Meystre, "Diffraction of ultra-cold fermions by a quantized light fields: Standing versus traveling waves," submitted to Phys. Rev. A.

"Bidirectional propagation of cold atoms in a "stadium" shaped magnetic guide", Saijun Wu, Wilbert Rooijakkers*, Pierre Striehl, Mara Prentiss, submitted to PRA

Y. Shin, M. Saba, A. Schirotzek, T. A. Pasquini, A. E. Leanhardt, D. E. Pritchard, and W. Ketterle: Distillation of Bose-Einstein condensates in a double-well potential. Preprint, cond-mat/ 0311514 (Phys. Rev. Lett. in print).

Ultracold Cs₂ Feshbach Spectroscopy, C. Chin, V. Vuletic, A.J. Kerman, S. Chu, E. Tiesinga, P.J. Leo, and C.J. Williams, submitted to Phys. Rev. A (December 2003).

Technical Reports submitted to ARO:

"Advanced matter wave sensors", Wolfgang Ketterle, Steven Chu, Eric Cornell and Carl Wieman (2002).

Participating Scientific Personnel

Steven Chu
Wolfgang Ketterle
Mark Kasevich
Dave Pritchard
Ewan Wright
Roy Glauber
Pierre Meystre
Mark Kasevich
Mara Prentiss

Report of Inventions

None